

## Study of $4n$ continuum in the $8\text{He} + 2\text{H}$ collisions at ACCULINNA-2 fragment separator

Recently in the experiment [M. Duer et al., Nature 606, 678 (2022)] a peak, reported as “resonance-like structure” in  $4n$  system, was observed in the  $1\text{H}(8\text{He}, p\alpha)4n$  reaction at  $E(4n) = 2.37$  MeV with  $\Gamma = 1.75$  MeV. Here we present the results of the experiment performed at ACCULINNA-2 fragment separator with a 26 AMeV secondary  $8\text{He}$  beam to study low-energy continuum of  $4n$  system in the reactions on deuterium target. These data were previously analyzed for the studies of  $7\text{H}$  and  $6\text{H}$  systems in the  $2\text{H}(8\text{He}, 3\text{He})7\text{H}$  and  $2\text{H}(8\text{He}, 4\text{He})6\text{H}$  reactions [I.A. Muzalevskii et al., Phys. Rev. C 103, 044313 (2021), E.Yu. Nikolskii et al., Phys. Rev. C 105, 064605 (2022)]. Evidence for a hump in the  $4n$  continuum at  $3.5 \pm 0.7$  and  $3.2 \pm 0.8$  MeV was observed in the  $2\text{H}(8\text{He}, 6\text{Li})4n$  and  $2\text{H}(8\text{He}, 3\text{He})7\text{H} \rightarrow 3\text{H} + 4n$  reactions, respectively. The obtained statistics is very low (6 and up to 40 events) corresponding to very low cross sections of few microbarns or tens of microbarns. The background conditions for the  $2\text{H}(8\text{He}, 6\text{Li})4n$  reaction are shown to be good, favoring the physical nature of the observed events. The  $2\text{H}(8\text{He}, 3\text{He})7\text{H} \rightarrow 3\text{H} + 4n$  process transforms to the  $2\text{H}(8\text{He}, 6\text{Li}^*)4n$  reaction in the limit of the highest  $7\text{H}$  decay energies. The population of the low-energy region in the  $4n$  spectrum is found to be correlated with the population of the lowest  $6\text{Li}$  states in the  $3\text{He} + 3\text{H}$  continuum. Theoretical calculations of  $8\text{He}$  in a five-body  $\alpha + 4n$  and of  $4n$  in a four-body hyperspherical models are presented. The  $8\text{He}$  wave function is shown to contain strong specific correlations, which may give rise to very low-energy structures in  $4n$  continuum in extreme-peripheral reaction scenarios.

### Section

Nuclear physics (Section 1)

**Primary author:** NIKOLSKII, Evgenii (NRC “Kurchatov Institute” / FLNR, JINR)

**Presenter:** NIKOLSKII, Evgenii (NRC “Kurchatov Institute” / FLNR, JINR)

**Track Classification:** The V International Scientific Forum “Nuclear Science and Technologies”: Nuclear physics (Section 1)